

**2-automatic sequences,
Lyapunov exponents and a
dynamical analogue of
Lehmer's Mahler measure problem**

Michael Coons

University of Newcastle, Australia

k-regular sequences

***k*-regular sequences**

Recall that an integer sequence \mathbf{f} is linear recurrent provided there is a $d \times d$ square matrix \mathbf{A} and $d \times 1$ vectors \mathbf{u} and \mathbf{v} such that

$$f(n) = \mathbf{u}^T \mathbf{A}^n \mathbf{v}.$$

In the generalisation, one doesn't consider the value of n specifically, but its base- k expansion $(n)_k = i_s \cdots i_0$.

k -regular sequences

Recall that an integer sequence \mathbf{f} is linear recurrent provided there is a $d \times d$ square matrix \mathbf{A} and $d \times 1$ vectors \mathbf{u} and \mathbf{v} such that

$$f(n) = \mathbf{u}^T \mathbf{A}^n \mathbf{v}.$$

In the generalisation, one doesn't consider the value of n specifically, but its base- k expansion $(n)_k = i_s \cdots i_0$.

In this setting, the sequence \mathbf{f} is k -regular if and only if there exist $d \times d$ square matrices $\mathbf{A}_0, \dots, \mathbf{A}_{k-1}$ and $d \times 1$ vectors \mathbf{u} and \mathbf{v} such that

$$f(n) = \mathbf{u}^T \mathbf{A}_w \mathbf{v} = \mathbf{u}^T \mathbf{A}_{i_s} \cdots \mathbf{A}_{i_0} \mathbf{v},$$

where $w = i_s \cdots i_0$ is the word corresponding to the base- k expansion.

Growth of k -regular sequences

Growth of k -regular sequences

We define the *growth exponent of f* , denoted $\text{GrExp}(f)$, by

$$\text{GrExp}(f) := \limsup_{\substack{n \rightarrow \infty \\ f(n) \neq 0}} \frac{\log |f(n)|}{\log n}.$$

The *joint spectral radius* of a finite set of matrices $\mathcal{A} = \{\mathbf{A}_0, \mathbf{A}_1, \dots, \mathbf{A}_{k-1}\}$, denoted $\rho(\mathcal{A})$, is defined as the real number

$$\rho(\mathcal{A}) = \limsup_{n \rightarrow \infty} \max_{0 \leq i_0, i_1, \dots, i_{n-1} \leq k-1} \left\| \mathbf{A}_{i_0} \mathbf{A}_{i_1} \cdots \mathbf{A}_{i_{n-1}} \right\|^{1/n},$$

where $\|\cdot\|$ is any (submultiplicative) matrix norm.

Growth of k -regular sequences

We define the *growth exponent of f* , denoted $\text{GrExp}(f)$, by

$$\text{GrExp}(f) := \limsup_{\substack{n \rightarrow \infty \\ f(n) \neq 0}} \frac{\log |f(n)|}{\log n}.$$

The *joint spectral radius* of a finite set of matrices $\mathcal{A} = \{\mathbf{A}_0, \mathbf{A}_1, \dots, \mathbf{A}_{k-1}\}$, denoted $\rho(\mathcal{A})$, is defined as the real number

$$\rho(\mathcal{A}) = \limsup_{n \rightarrow \infty} \max_{0 \leq i_0, i_1, \dots, i_{n-1} \leq k-1} \left\| \mathbf{A}_{i_0} \mathbf{A}_{i_1} \cdots \mathbf{A}_{i_{n-1}} \right\|^{1/n},$$

where $\|\cdot\|$ is any (submultiplicative) matrix norm.

Theorem. *Let $k \geq 1$ and $d \geq 1$ be integers and $f : \mathbb{Z}_{\geq 0} \rightarrow \mathbb{K}$ be a (not eventually zero) k -regular sequence. If \mathcal{A}_f is any collection of k integer matrices associated to a basis of the \mathbb{K} -vector space $\langle \text{Ker}_k(f) \rangle_{\mathbb{K}}$, then*

$$\log_k \rho(\mathcal{A}_f) = \text{GrExp}(f),$$

where \log_k denotes the base- k logarithm.

An example: Stern's sequence

An example: Stern's sequence

For example, the *Stern sequence*, $\{s(n)\}_{n \geq 0}$ is given by $s(0) = 0$, $s(1) = 1$, and when $n \geq 1$, by the pair of recurrences

$$s(2n) = s(n) \quad \text{and} \quad s(2n + 1) = s(n) + s(n + 1).$$

These recurrences are determined by the matrix representation of the Stern sequence, where

$$\{\mathbf{A}_0, \mathbf{A}_1\} = \left\{ \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} \right\}.$$

An example: Stern's sequence

For example, the *Stern sequence*, $\{s(n)\}_{n \geq 0}$ is given by $s(0) = 0$, $s(1) = 1$, and when $n \geq 1$, by the pair of recurrences


$$s(2n) = s(n) \quad \text{and} \quad s(2n + 1) = s(n) + s(n + 1).$$

These recurrences are determined by the matrix representation of the Stern sequence, where

$$\{\mathbf{A}_0, \mathbf{A}_1\} = \left\{ \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} \right\}.$$

THEOREM 1.1. *Let $\{a(n)\}_{n \geq 0}$ denote the Stern sequence. Then*

$$\limsup_{n \rightarrow \infty} \frac{a(n)}{n^{\log_2 \varphi}} = \frac{\varphi}{\sqrt{5}} \left(\frac{3}{2} \right)^{\log_2 \varphi} = \frac{\varphi^{\log_2 3}}{\sqrt{5}} = 0.9588541900 \dots$$



Regular to Mahler

Regular to Mahler

Definition. A function $F(z) \in \mathbb{Z}[z]$ is called a *k-Mahler function* provided there are integers $k \geq 2$ and $d \geq 1$ such that

$$a_0(z)F(z) + a_1(z)F(z^k) + \cdots + a_d(z)F(z^{k^d}) = 0,$$

for some polynomials $a_0(z), \dots, a_d(z) \in \mathbb{Z}[z]$.

Regular to Mahler

Definition. A function $F(z) \in \mathbb{Z}[z]$ is called a *k-Mahler function* provided there are integers $k \geq 2$ and $d \geq 1$ such that

$$a_0(z)F(z) + a_1(z)F(z^k) + \cdots + a_d(z)F(z^{k^d}) = 0,$$

for some polynomials $a_0(z), \dots, a_d(z) \in \mathbb{Z}[z]$.

Theorem. (Becker) *If $f(n)$ is a k-regular sequence, then $\sum_{n \geq 0} f(n)z^n$ is a k-Mahler function.*

Regular to Mahler

Definition. A function $F(z) \in \mathbb{Z}[z]$ is called a *k-Mahler function* provided there are integers $k \geq 2$ and $d \geq 1$ such that

$$a_0(z)F(z) + a_1(z)F(z^k) + \cdots + a_d(z)F(z^{k^d}) = 0,$$

for some polynomials $a_0(z), \dots, a_d(z) \in \mathbb{Z}[z]$.

Theorem. (Becker) *If $f(n)$ is a k-regular sequence, then $\sum_{n \geq 0} f(n)z^n$ is a k-Mahler function.*

The function $S(z) = (1/z) \sum_{n \geq 0} s(n)z^n$ satisfies the 2-Mahler equation

$$S(z) = (1 + z + z^2)S(z^2).$$

A transcendence test for Mahler functions

Let $k \geq 2$ and $d \geq 1$ be integers and $F(z)$ be a k -Mahler function converging inside the unit disc satisfying

$$a_0(z)F(z) + a_1(z)F(z^k) + \cdots + a_d(z)F(z^{k^d}) = 0,$$

for polynomials $a_0(z), \dots, a_d(z) \in \mathbb{C}[z]$. Set $a_i := a_i(1)$ and form the polynomial

$$p_F(\lambda) := a_0\lambda^d + a_1\lambda^{d-1} + \cdots + a_{d-1}\lambda + a_d.$$

If $a_0a_d \neq 0$ and $p_F(\lambda)$ has distinct roots, then the function $F(z)$ is transcendental over $\mathbb{C}(z)$ provided

- $p(k^n) \neq 0$ for all $n \in \mathbb{Z}$ or
- the eigenvalue $\lambda_F \neq k^n$ for any $n \in \mathbb{Z}$.

If $\lambda_F = k^n$ for some $n \in \mathbb{Z}$, the test is inconclusive.

Radial asymptotics of Mahler functions

Radial asymptotics of Mahler functions

Theorem. *Let $F(z)$ be a k -Mahler function whose characteristic polynomial $p_F(\lambda)$ has distinct roots. Then there is an eigenvalue λ_F with $p_F(\lambda_F) = 0$, such that as $z \rightarrow 1^-$*

$$F(z) = \frac{C(z)}{(1-z)^{\log_k \lambda_F}} (1 + o(1)),$$

where \log_k denotes the principal value of the base- k logarithm and $C(z)$ is a real-analytic nonzero oscillatory term, which on the interval $(0, 1)$ is bounded away from 0 and ∞ , and satisfies $C(z) = C(z^k)$.

Questions about the asymptotic behaviour of Mahler functions are quite classical, and some special cases of the theorem above are known. See, e.g., Mahler (1940), de Bruijn (1948), Dumas (1993), Dumas and Flajolet (1996), and most recently Brent, Coons, and Zudilin (2015).

Properties via matrix cocycles

Properties via matrix cocycles

Mahler functions are coordinates of a vector $\mathbf{F}(z) := [F_1(z), \dots, F_d(z)]^T$ such that there is a matrix of rational functions $\mathbf{A}(z)$ such that

$$\mathbf{F}(z) = \mathbf{A}(z)\mathbf{F}(z^k).$$

Properties via matrix cocycles

Mahler functions are coordinates of a vector $\mathbf{F}(z) := [F_1(z), \dots, F_d(z)]^T$ such that there is a matrix of rational functions $\mathbf{A}(z)$ such that

$$\mathbf{F}(z) = \mathbf{A}(z)\mathbf{F}(z^k).$$

For every n , we have $\mathbf{F}(z) = \underbrace{\mathbf{A}(z)\mathbf{A}(z^k) \cdots \mathbf{A}(z^{k^{n-1}})}_{\text{matrix cocycle}} \mathbf{F}(z^{k^n})$.

Properties via matrix cocycles

Mahler functions are coordinates of a vector $\mathbf{F}(z) := [F_1(z), \dots, F_d(z)]^T$ such that there is a matrix of rational functions $\mathbf{A}(z)$ such that

$$\mathbf{F}(z) = \mathbf{A}(z)\mathbf{F}(z^k).$$

For every n , we have $\mathbf{F}(z) = \underbrace{\mathbf{A}(z)\mathbf{A}(z^k) \cdots \mathbf{A}(z^{k^{n-1}})}_{\text{matrix cocycle}} \mathbf{F}(z^{k^n})$.

Properties of this cocycle can give you:

- transcendence results
- algebraic independence results
- irrationality measures

Approximation by rationals

Approximation by rationals

Definition. Let ξ be a real number. The *irrationality exponent* $\mu(\xi)$ is defined as the supremum of the set of real numbers μ such that the inequality

$$\left| \xi - \frac{p}{q} \right| < \frac{1}{q^\mu}$$

has infinitely many solutions $(p, q) \in \mathbb{Z} \times \mathbb{N}$.

Approximation by rationals

Definition. Let ξ be a real number. The *irrationality exponent* $\mu(\xi)$ is defined as the supremum of the set of real numbers μ such that the inequality

$$\left| \xi - \frac{p}{q} \right| < \frac{1}{q^\mu}$$

has infinitely many solutions $(p, q) \in \mathbb{Z} \times \mathbb{N}$.

Theorem (Dirichlet 1842). *Let ξ be a real number. If $\mu(\xi) < 2$, then $\mu(\xi) = 1$. Moreover, all number ξ with $\mu(\xi) = 1$ are rational.*

Approximation by rationals

Definition. Let ξ be a real number. The *irrationality exponent* $\mu(\xi)$ is defined as the supremum of the set of real numbers μ such that the inequality

$$\left| \xi - \frac{p}{q} \right| < \frac{1}{q^\mu}$$

has infinitely many solutions $(p, q) \in \mathbb{Z} \times \mathbb{N}$.

Theorem (Dirichlet 1842). *Let ξ be a real number. If $\mu(\xi) < 2$, then $\mu(\xi) = 1$. Moreover, all number ξ with $\mu(\xi) = 1$ are rational.*

Theorem (Liouville 1844). *If ξ is algebraic of degree d , then $\mu(\xi) \leq d$.*

Approximation by rationals

Definition. Let ξ be a real number. The *irrationality exponent* $\mu(\xi)$ is defined as the supremum of the set of real numbers μ such that the inequality

$$\left| \xi - \frac{p}{q} \right| < \frac{1}{q^\mu}$$

has infinitely many solutions $(p, q) \in \mathbb{Z} \times \mathbb{N}$.

Theorem (Dirichlet 1842). *Let ξ be a real number. If $\mu(\xi) < 2$, then $\mu(\xi) = 1$. Moreover, all number ξ with $\mu(\xi) = 1$ are rational.*

Theorem (Liouville 1844). *If ξ is algebraic of degree d , then $\mu(\xi) \leq d$.*

THEOREM 1.1. *A Mahler number cannot be a Liouville number.*

Mahler measure and possible gaps

Mahler measure and possible gaps

Definition. Let $p(z) = a_0 \prod_{i=0}^s (z - \alpha_i) \in \mathbb{C}[z]$. The (logarithmic) Mahler measure of $p(z)$ is given by

$$\mathfrak{m}(p) := \log |a_0| + \sum_{i=0}^s \log(\max\{|\alpha_i|, 1\}).$$

Mahler measure and possible gaps

Definition. Let $p(z) = a_0 \prod_{i=0}^s (z - \alpha_i) \in \mathbb{C}[z]$. The (logarithmic) Mahler measure of $p(z)$ is given by

$$\mathfrak{m}(p) := \log |a_0| + \sum_{i=0}^s \log(\max\{|\alpha_i|, 1\}).$$

Note also Jensen's formula $\mathfrak{m}(p) := \int_0^1 \log |p(e^{2\pi it})| dt$.

Mahler measure and possible gaps

Definition. Let $p(z) = a_0 \prod_{i=0}^s (z - \alpha_i) \in \mathbb{C}[z]$. The (logarithmic) Mahler measure of $p(z)$ is given by

$$\mathfrak{m}(p) := \log |a_0| + \sum_{i=0}^s \log(\max\{|\alpha_i|, 1\}).$$

Note also Jensen's formula $\mathfrak{m}(p) := \int_0^1 \log |p(e^{2\pi it})| dt$.

Lehmer found $\mathfrak{m}(1 + z - z^3 - z^4 - z^5 - z^6 - z^7 + z^9 + z^{10}) \approx \log(1.176281)$.

Mahler measure and possible gaps

Definition. Let $p(z) = a_0 \prod_{i=0}^s (z - \alpha_i) \in \mathbb{C}[z]$. The (logarithmic) Mahler measure of $p(z)$ is given by

$$\mathfrak{m}(p) := \log |a_0| + \sum_{i=0}^s \log(\max\{|\alpha_i|, 1\}).$$

Note also Jensen's formula $\mathfrak{m}(p) := \int_0^1 \log |p(e^{2\pi it})| dt$.

Lehmer found $\mathfrak{m}(1 + z - z^3 - z^4 - z^5 - z^6 - z^7 + z^9 + z^{10}) \approx \log(1.176281)$.

Lehmer's problem. Does there exist a constant $c > 0$ such that any irreducible non-cyclotomic polynomial p with integer coefficients satisfies $\mathfrak{m}(p) \geq c$?

Boyd's observation and height-one polynomials

Boyd's observation and height-one polynomials

Lehmer's problem. Does there exist a constant $c > 0$ such that any irreducible non-cyclotomic polynomial p with integer coefficients satisfies $\mathfrak{m}(p) \geq c$?

Boyd's observation and height-one polynomials

Lehmer's problem. Does there exist a constant $c > 0$ such that any irreducible non-cyclotomic polynomial p with integer coefficients satisfies $\mathfrak{m}(p) \geq c$?

For any integer polynomial p with $\mathfrak{m}(p) < \log(2)$ there is an integer polynomial q such that pq has height 1.

Boyd's observation and height-one polynomials

Lehmer's problem. Does there exist a constant $c > 0$ such that any irreducible non-cyclotomic polynomial p with integer coefficients satisfies $\mathfrak{m}(p) \geq c$?

For any integer polynomial p with $\mathfrak{m}(p) < \log(2)$ there is an integer polynomial q such that pq has height 1.

Boyd observed that, in his experience, such a q can be taken to be cyclotomic and of fairly small degree relative to the degree of p .

Boyd's observation and height-one polynomials

Lehmer's problem. Does there exist a constant $c > 0$ such that any irreducible non-cyclotomic polynomial p with integer coefficients satisfies $\mathfrak{m}(p) \geq c$?

For any integer polynomial p with $\mathfrak{m}(p) < \log(2)$ there is an integer polynomial q such that pq has height 1.

Boyd observed that, in his experience, such a q can be taken to be cyclotomic and of fairly small degree relative to the degree of p .

In **this talk** I want to show you that Lehmer's problem, restricted to height-one polynomials, is a property of a matrix cocycle associated to a 2-automatic sequence.

Fourier matrix of a 2-automatic sequence

Fourier matrix of a 2-automatic sequence

Definition. A 2-automatic sequence ϱ is defined on $\Sigma_2 := \{0, 1\}$ by

$$\varrho : \begin{cases} 0 \mapsto w_0 \\ 1 \mapsto w_1, \end{cases}$$

where w_0 and w_1 are finite words over Σ_2 of equal length $|w_0| = |w_1| = L \geq 2$.

Fourier matrix of a 2-automatic sequence

Definition. A 2-automatic sequence ϱ is defined on $\Sigma_2 := \{0, 1\}$ by

$$\varrho : \begin{cases} 0 \mapsto w_0 \\ 1 \mapsto w_1, \end{cases}$$

where w_0 and w_1 are finite words over Σ_2 of equal length $|w_0| = |w_1| = L \geq 2$.

All sequences will be assumed to be *primitive* and *aperiodic*.

Fourier matrix of a 2-automatic sequence

Definition. A 2-automatic sequence ϱ is defined on $\Sigma_2 := \{0, 1\}$ by

$$\varrho : \begin{cases} 0 \mapsto w_0 \\ 1 \mapsto w_1, \end{cases}$$

where w_0 and w_1 are finite words over Σ_2 of equal length $|w_0| = |w_1| = L \geq 2$.

All sequences will be assumed to be *primitive* and *aperiodic*.

For $0 \leq i, j \leq 1$, let T_{ij} be the set of all positions m where the letter i appears in w_j , and let $T := (T_{ij})_{0 \leq i, j \leq 1}$ be the resulting 2×2 -matrix.

Fourier matrix of a 2-automatic sequence

Definition. A 2-automatic sequence ϱ is defined on $\Sigma_2 := \{0, 1\}$ by

$$\varrho : \begin{cases} 0 \mapsto w_0 \\ 1 \mapsto w_1, \end{cases}$$

where w_0 and w_1 are finite words over Σ_2 of equal length $|w_0| = |w_1| = L \geq 2$.

All sequences will be assumed to be *primitive* and *aperiodic*.

For $0 \leq i, j \leq 1$, let T_{ij} be the set of all positions m where the letter i appears in w_j , and let $T := (T_{ij})_{0 \leq i, j \leq 1}$ be the resulting 2×2 -matrix.

Note that the substitution matrix M_ϱ , satisfies $M_\varrho = (\text{card}(T_{ij}))_{0 \leq i, j \leq 1}$.

Fourier matrix of a 2-automatic sequence

Definition. A 2-automatic sequence ϱ is defined on $\Sigma_2 := \{0, 1\}$ by

$$\varrho : \begin{cases} 0 \mapsto w_0 \\ 1 \mapsto w_1, \end{cases}$$

where w_0 and w_1 are finite words over Σ_2 of equal length $|w_0| = |w_1| = L \geq 2$.

All sequences will be assumed to be *primitive* and *aperiodic*.

For $0 \leq i, j \leq 1$, let T_{ij} be the set of all positions m where the letter i appears in w_j , and let $T := (T_{ij})_{0 \leq i, j \leq 1}$ be the resulting 2×2 -matrix.

Note that the substitution matrix M_ϱ , satisfies $M_\varrho = (\text{card}(T_{ij}))_{0 \leq i, j \leq 1}$.

Definition. Using T , we build a matrix of pure point measures $\delta_T = (\delta_{T_{ij}})_{0 \leq i, j \leq 1}$, where $\delta_S := \sum_{x \in S} \delta_x$ with $\delta_\emptyset = 0$. This gives rise to an analytic matrix-valued function via

$$B(k) := \overline{\widehat{\delta_T}(k)},$$

which we call the *Fourier matrix* of ϱ .

Two paradigmatic examples

Two paradigmatic examples

Example. Consider the Thue–Morse substitution, as given by

$$\varrho_{\text{TM}} : \begin{cases} 0 \mapsto 01 \\ 1 \mapsto 10. \end{cases}$$

Here, one has $T_{\text{TM}} = \begin{pmatrix} \{0\} & \{1\} \\ \{1\} & \{0\} \end{pmatrix}$, which gives

$$\delta_{T_{\text{TM}}} = \begin{pmatrix} \delta_0 & \delta_1 \\ \delta_1 & \delta_0 \end{pmatrix} \quad \text{and} \quad B_{\text{TM}}(k) = \begin{pmatrix} 1 & e^{2\pi i k} \\ e^{2\pi i k} & 1 \end{pmatrix}.$$

Two paradigmatic examples

Example. Consider the Thue–Morse substitution, as given by

$$\varrho_{\text{TM}} : \begin{cases} 0 \mapsto 01 \\ 1 \mapsto 10. \end{cases}$$

Here, one has $T_{\text{TM}} = \begin{pmatrix} \{0\} & \{1\} \\ \{1\} & \{0\} \end{pmatrix}$, which gives

$$\delta_{T_{\text{TM}}} = \begin{pmatrix} \delta_0 & \delta_1 \\ \delta_1 & \delta_0 \end{pmatrix} \quad \text{and} \quad B_{\text{TM}}(k) = \begin{pmatrix} 1 & e^{2\pi i k} \\ e^{2\pi i k} & 1 \end{pmatrix}.$$

Example. On the other hand, for the period doubling substitution,

$$\varrho_{\text{pd}} : \begin{cases} 0 \mapsto 01 \\ 1 \mapsto 00, \end{cases}$$

the corresponding matrices are $T_{\text{pd}} = \begin{pmatrix} \{0\} & \{0,1\} \\ \{1\} & \emptyset \end{pmatrix}$ together with

$$\delta_{T_{\text{pd}}} = \begin{pmatrix} \delta_0 & \delta_0 + \delta_1 \\ \delta_1 & 0 \end{pmatrix} \quad \text{and} \quad B_{\text{pd}}(k) = \begin{pmatrix} 1 & 1 + e^{2\pi i k} \\ e^{2\pi i k} & 0 \end{pmatrix}.$$

Lyapunov exponents and Fourier cocycles

Lyapunov exponents and Fourier cocycles

We use the ergodic transformation $k \mapsto Lk \bmod 1$ defined on the 1-torus and the Fourier matrix to build the matrix cocycle

$$B^{(n)}(k) := B(k)B(Lk) \cdots B(L^{n-1}k).$$

Lyapunov exponents and Fourier cocycles

We use the ergodic transformation $k \mapsto Lk \bmod 1$ defined on the 1-torus and the Fourier matrix to build the matrix cocycle

$$B^{(n)}(k) := B(k)B(Lk) \cdots B(L^{n-1}k).$$

The inverse cocycle $(B^{(n)}(k))^{-1}$ exists for almost all $k \in \mathbb{R}$, because $\det B(k) = 0$ for at most a countable subset of \mathbb{R} .

Lyapunov exponents and Fourier cocycles

We use the ergodic transformation $k \mapsto Lk \bmod 1$ defined on the 1-torus and the Fourier matrix to build the matrix cocycle

$$B^{(n)}(k) := B(k)B(Lk) \cdots B(L^{n-1}k).$$

The inverse cocycle $(B^{(n)}(k))^{-1}$ exists for almost all $k \in \mathbb{R}$, because $\det B(k) = 0$ for at most a countable subset of \mathbb{R} .

- If $v \in \mathbb{C}^2$ is any (fixed) row vector, the values

$$\chi^B(v, k) := \lim_{n \rightarrow \infty} \frac{1}{n} \log \|v B^{(n)}(k)\|$$

exist for almost every $k \in \mathbb{R}$ and are constant on a set of full measure.

Lyapunov exponents and Fourier cocycles

We use the ergodic transformation $k \mapsto Lk \bmod 1$ defined on the 1-torus and the Fourier matrix to build the matrix cocycle

$$B^{(n)}(k) := B(k)B(Lk) \cdots B(L^{n-1}k).$$

The inverse cocycle $(B^{(n)}(k))^{-1}$ exists for almost all $k \in \mathbb{R}$, because $\det B(k) = 0$ for at most a countable subset of \mathbb{R} .

- If $v \in \mathbb{C}^2$ is any (fixed) row vector, the values

$$\chi^B(v, k) := \lim_{n \rightarrow \infty} \frac{1}{n} \log \|v B^{(n)}(k)\|$$

exist for almost every $k \in \mathbb{R}$ and are constant on a set of full measure.

- As a function of v , they take at most two values in this case.

Lyapunov exponents and Fourier cocycles

We use the ergodic transformation $k \mapsto Lk \bmod 1$ defined on the 1-torus and the Fourier matrix to build the matrix cocycle

$$B^{(n)}(k) := B(k)B(Lk) \cdots B(L^{n-1}k).$$

The inverse cocycle $(B^{(n)}(k))^{-1}$ exists for almost all $k \in \mathbb{R}$, because $\det B(k) = 0$ for at most a countable subset of \mathbb{R} .

- If $v \in \mathbb{C}^2$ is any (fixed) row vector, the values

$$\chi^B(v, k) := \lim_{n \rightarrow \infty} \frac{1}{n} \log \|vB^{(n)}(k)\|$$

exist for almost every $k \in \mathbb{R}$ and are constant on a set of full measure.

- As a function of v , they take at most two values in this case.
- A vector v from the Oseledec subspace $\mathcal{V}_{i+1} \setminus \mathcal{V}_i$ satisfies the property that, for almost every $k \in \mathbb{R}$, the norm $\|vB^{(n)}(k)\|$ grows like $e^{n\chi_{i+1}^B}$ as $n \rightarrow \infty$.

- For invertible cocycles, these exponents have v -independent forms

$$\chi_{\max}^B(k) := \lim_{n \rightarrow \infty} \frac{1}{n} \log \|B^{(n)}(k)\| \quad \text{and}$$
$$\chi_{\min}^B(k) := - \lim_{n \rightarrow \infty} \frac{1}{n} \log \|(B^{(n)}(k))^{-1}\|.$$

- For invertible cocycles, these exponents have v -independent forms

$$\chi_{\max}^B(k) := \lim_{n \rightarrow \infty} \frac{1}{n} \log \|B^{(n)}(k)\| \quad \text{and}$$
$$\chi_{\min}^B(k) := - \lim_{n \rightarrow \infty} \frac{1}{n} \log \|(B^{(n)}(k))^{-1}\|.$$

- These values are independent of the norm.

- For invertible cocycles, these exponents have v -independent forms

$$\chi_{\max}^B(k) := \lim_{n \rightarrow \infty} \frac{1}{n} \log \|B^{(n)}(k)\| \quad \text{and}$$
$$\chi_{\min}^B(k) := - \lim_{n \rightarrow \infty} \frac{1}{n} \log \|(B^{(n)}(k))^{-1}\|.$$

- These values are independent of the norm.
- This means that we are dealing with two *numbers*, χ_{\max}^B and χ_{\min}^B .

- For invertible cocycles, these exponents have v -independent forms

$$\chi_{\max}^B(k) := \lim_{n \rightarrow \infty} \frac{1}{n} \log \|B^{(n)}(k)\| \quad \text{and}$$

$$\chi_{\min}^B(k) := - \lim_{n \rightarrow \infty} \frac{1}{n} \log \|(B^{(n)}(k))^{-1}\|.$$

- These values are independent of the norm.
- This means that we are dealing with two *numbers*, χ_{\max}^B and χ_{\min}^B .
- Lyapunov regularity guarantees for almost every $k \in \mathbb{R}$ that

$$\chi_{\min}^B(k) + \chi_{\max}^B(k) = \lim_{n \rightarrow \infty} \frac{1}{n} \log |\det(B^{(n)}(k))|.$$

- For invertible cocycles, these exponents have v -independent forms

$$\chi_{\max}^B(k) := \lim_{n \rightarrow \infty} \frac{1}{n} \log \|B^{(n)}(k)\| \quad \text{and}$$

$$\chi_{\min}^B(k) := - \lim_{n \rightarrow \infty} \frac{1}{n} \log \| (B^{(n)}(k))^{-1} \|.$$

- These values are independent of the norm.
- This means that we are dealing with two *numbers*, χ_{\max}^B and χ_{\min}^B .
- Lyapunov regularity guarantees for almost every $k \in \mathbb{R}$ that

$$\chi_{\min}^B(k) + \chi_{\max}^B(k) = \lim_{n \rightarrow \infty} \frac{1}{n} \log |\det(B^{(n)}(k))|.$$

Theorem (Baake, C. and Mañibo). *For any primitive, 2-automatic sequence ϱ , the extremal Lyapunov exponents are explicitly given by*

$$\chi_{\min}^B = 0 \quad \text{and} \quad \chi_{\max}^B = \mathfrak{m}(Q - R),$$

with Q and R defined in the next lemma.

Lemma. Consider the sets

$$P_a := \{m \mid \mathcal{C}_m = \begin{bmatrix} 0 \\ 1 \end{bmatrix}\} \quad \text{and} \quad P_b := \{m \mid \mathcal{C}_m = \begin{bmatrix} 1 \\ 0 \end{bmatrix}\},$$

which collect bijective positions of equal type. Further, let $z = e^{2\pi i k}$ and set

$$Q(z) := \overline{\widehat{\delta}_{P_a}(k)} \quad \text{and} \quad R(z) := \overline{\widehat{\delta}_{P_b}(k)}.$$

Then, $\det(B(k)) = p_L(z) \cdot (Q - R)(z)$, where $p_L(z) = 1 + z + \dots + z^{L-1}$.

Lemma. Consider the sets

$$P_a := \{m \mid \mathcal{C}_m = \begin{bmatrix} 0 \\ 1 \end{bmatrix}\} \quad \text{and} \quad P_b := \{m \mid \mathcal{C}_m = \begin{bmatrix} 1 \\ 0 \end{bmatrix}\},$$

which collect bijective positions of equal type. Further, let $z = e^{2\pi ik}$ and set

$$Q(z) := \overline{\widehat{\delta}_{P_a}(k)} \quad \text{and} \quad R(z) := \overline{\widehat{\delta}_{P_b}(k)}.$$

Then, $\det(B(k)) = p_L(z) \cdot (Q - R)(z)$, where $p_L(z) = 1 + z + \dots + z^{L-1}$.

Proof. Similar to the definitions of Q and R above, define $P_0 := \{m \mid \mathcal{C}_m = \begin{bmatrix} 0 \\ 0 \end{bmatrix}\}$ and $P_1 := \{m \mid \mathcal{C}_m = \begin{bmatrix} 1 \\ 1 \end{bmatrix}\}$, and let

$$S_0(z) := \overline{\widehat{\delta}_{P_0}(k)} \quad \text{and} \quad S_1(z) := \overline{\widehat{\delta}_{P_1}(k)}.$$

In general, the Fourier matrix of ϱ satisfies

$$B(k) = \begin{pmatrix} (S_0 + Q)(z) & (S_0 + R)(z) \\ (S_1 + R)(z) & (S_1 + Q)(z) \end{pmatrix} \quad \text{with } z = e^{2\pi ik}.$$

Since there are only four distinct column types, we see that $S_0 + S_1 + Q + R = p_L$. One can now verify the lemma by direct computation. \square

Main result

Main result

Theorem (Baake, C. and Mañibo). *For any primitive, 2-automatic sequence ϱ , the extremal Lyapunov exponents are explicitly given by*

$$\chi_{\min}^B = 0 \quad \text{and} \quad \chi_{\max}^B = \mathfrak{m}(Q - R),$$

with Q and R defined in the previous lemma.

Main result

Theorem (Baake, C. and Mañibo). *For any primitive, 2-automatic sequence ϱ , the extremal Lyapunov exponents are explicitly given by*

$$\chi_{\min}^B = 0 \quad \text{and} \quad \chi_{\max}^B = \mathfrak{m}(Q - R),$$

with Q and R defined in the previous lemma.

Proof. By the ergodic theorem, multiplicativity of the determinant, and additivity of the logarithm, for almost every $k \in \mathbb{R}$

$$\begin{aligned} \chi_{\min}^B(k) + \chi_{\max}^B(k) &= \lim_{n \rightarrow \infty} \frac{1}{n} \log |\det(B^{(n)}(k))| \\ &= \int_0^1 \log |(Q - R)(e^{2\pi it})| dt = \mathfrak{m}(Q - R). \end{aligned}$$

Main result

Theorem (Baake, C. and Mañibo). *For any primitive, 2-automatic sequence ϱ , the extremal Lyapunov exponents are explicitly given by*

$$\chi_{\min}^B = 0 \quad \text{and} \quad \chi_{\max}^B = \mathfrak{m}(Q - R),$$

with Q and R defined in the previous lemma.

Proof. By the ergodic theorem, multiplicativity of the determinant, and additivity of the logarithm, for almost every $k \in \mathbb{R}$

$$\begin{aligned} \chi_{\min}^B(k) + \chi_{\max}^B(k) &= \lim_{n \rightarrow \infty} \frac{1}{n} \log |\det(B^{(n)}(k))| \\ &= \int_0^1 \log |(Q - R)(e^{2\pi it})| dt = \mathfrak{m}(Q - R). \end{aligned}$$

The row vector $(1, 1)$ is a left eigenvector of $B(k)$, for all k , with eigenvalue $p_L(e^{2\pi ik})$. Hence, we get one of the exponents to be $\chi_1^B = \mathfrak{m}(p_L) = 0$. From the sum above, and from the non-negativity of the Mahler measure, we then get that the exponent corresponding to this invariant subspace is the minimal one, $\chi_1^B = \chi_{\min}^B$, the other being $\chi_{\max}^B = \mathfrak{m}(Q - R)$. \square

Important point

Important point

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log \left| \det \underbrace{\left(B(k) B(Lk) \cdots B(L^{n-1}k) \right)}_{\text{matrix cocycle}} \right| = \mathfrak{m}(Q - R).$$

Important point

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log \left| \det \underbrace{\left(B(k) B(Lk) \cdots B(L^{n-1}k) \right)}_{\text{matrix cocycle}} \right| = \mathfrak{m}(Q - R).$$

There is a massive amount of structure to the matrix $B(k)$.

Important point

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log \left| \det \underbrace{\left(B(k) B(Lk) \cdots B(L^{n-1}k) \right)}_{\text{matrix cocycle}} \right| = \mathfrak{m}(Q - R).$$

There is a massive amount of structure to the matrix $B(k)$.

$$B(k) = \begin{pmatrix} (S_0 + Q)(z) & (S_0 + R)(z) \\ (S_1 + R)(z) & (S_1 + Q)(z) \end{pmatrix} \quad \text{with } z = e^{2\pi i k}.$$

Important point

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log \left| \det \underbrace{\left(B(k) B(Lk) \cdots B(L^{n-1}k) \right)}_{\text{matrix cocycle}} \right| = \mathfrak{m}(Q - R).$$

There is a massive amount of structure to the matrix $B(k)$.

$$B(k) = \begin{pmatrix} (S_0 + Q)(z) & (S_0 + R)(z) \\ (S_1 + R)(z) & (S_1 + Q)(z) \end{pmatrix} \quad \text{with } z = e^{2\pi i k}.$$

- One can easily bound heights
- bound degrees of the entries,
- and see how they are related.

Example: Littlewood polynomials and bijective morphisms

Example: Littlewood polynomials and bijective morphisms

Definition. A polynomial $q(z) = \sum_{m=0}^{n-1} c_m z^m$ of degree $n - 1$ with coefficients $c_m \in \{-1, 1\}$ is called a *Littlewood polynomial of order $n - 1$* .

Example: Littlewood polynomials and bijective morphisms

Definition. A polynomial $q(z) = \sum_{m=0}^{n-1} c_m z^m$ of degree $n - 1$ with coefficients $c_m \in \{-1, 1\}$ is called a *Littlewood polynomial of order $n - 1$* .

Let \mathcal{C}_m be the m^{th} column of ρ , and starting with $q(z)$, we choose \mathcal{C}_m to be

$$\mathcal{C}_m = \begin{cases} \begin{bmatrix} 0 \\ 1 \end{bmatrix}, & \text{if } c_m = 1, \\ \begin{bmatrix} 1 \\ 0 \end{bmatrix}, & \text{if } c_m = -1, \end{cases}$$

and we build the morphism by looking at the concatenation $\mathcal{C}_0 \mathcal{C}_1 \cdots \mathcal{C}_{L-1}$.

Example: Littlewood polynomials and bijective morphisms

Definition. A polynomial $q(z) = \sum_{m=0}^{n-1} c_m z^m$ of degree $n - 1$ with coefficients $c_m \in \{-1, 1\}$ is called a *Littlewood polynomial of order $n - 1$* .

Let \mathcal{C}_m be the m^{th} column of ϱ , and starting with $q(z)$, we choose \mathcal{C}_m to be

$$\mathcal{C}_m = \begin{cases} \begin{bmatrix} 0 \\ 1 \end{bmatrix}, & \text{if } c_m = 1, \\ \begin{bmatrix} 1 \\ 0 \end{bmatrix}, & \text{if } c_m = -1, \end{cases}$$

and we build the morphism by looking at the concatenation $\mathcal{C}_0 \mathcal{C}_1 \cdots \mathcal{C}_{L-1}$.

There are only two possible column types present, so $P_a \cup P_b = \{0, 1, \dots, L-1\}$, and also that the resulting morphism ϱ must be bijective.

Example: Littlewood polynomials and bijective morphisms

Definition. A polynomial $q(z) = \sum_{m=0}^{n-1} c_m z^m$ of degree $n - 1$ with coefficients $c_m \in \{-1, 1\}$ is called a *Littlewood polynomial of order $n - 1$* .

Let \mathcal{C}_m be the m^{th} column of ϱ , and starting with $q(z)$, we choose \mathcal{C}_m to be

$$\mathcal{C}_m = \begin{cases} \begin{bmatrix} 0 \\ 1 \end{bmatrix}, & \text{if } c_m = 1, \\ \begin{bmatrix} 1 \\ 0 \end{bmatrix}, & \text{if } c_m = -1, \end{cases}$$

and we build the morphism by looking at the concatenation $\mathcal{C}_0 \mathcal{C}_1 \cdots \mathcal{C}_{L-1}$.

There are only two possible column types present, so $P_a \cup P_b = \{0, 1, \dots, L-1\}$, and also that the resulting morphism ϱ must be bijective.

By construction, we have

$$\chi_{\max}^B = \mathfrak{m}(Q - R) = \mathfrak{m}(q) \quad \text{and} \quad B(k) = \begin{pmatrix} Q(z) & R(z) \\ R(z) & Q(z) \end{pmatrix} \quad \text{where } z = e^{2\pi i k}.$$

The substitution corresponding to $q = Q - R$ is essentially unique, up to the obvious freedom that emerges from the relation $\mathbf{m}(-q) = \mathbf{m}(q)$, that is, from changing all signs. This is the case precisely because a given sequence of signs uniquely specifies the columns of ϱ . For example, let us consider the polynomial

$$q(z) = -1 - z + z^2 - z^3 + z^4,$$

whence we get the substitutions

$$\varrho_q : \begin{cases} 0 \mapsto 11010 \\ 1 \mapsto 00101 \end{cases} \quad \text{and} \quad \varrho_{-q} : \begin{cases} 0 \mapsto 00101 \\ 1 \mapsto 11010 \end{cases}$$

with associated Fourier matrices

$$B_q(k) = \begin{pmatrix} e^{4\pi ik} + e^{8\pi ik} & 1 + e^{2\pi ik} + e^{6\pi ik} \\ 1 + e^{2\pi ik} + e^{6\pi ik} & e^{4\pi ik} + e^{8\pi ik} \end{pmatrix} \quad \text{and} \quad B_{-q}(k) = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} B_q(k).$$

Both induce a cocycle whose maximum Lyapunov exponent is

$$\chi_{\max}^B = \mathbf{m}(q) \approx 0.656256$$

Example: Newman polynomials and choices

Example: Newman polynomials and choices

For the class of $\{0, 1\}$ -polynomials, also known as *Newman polynomials*, one has $R = 0$. The associated Fourier matrix is

$$B(k) = \begin{pmatrix} (S_0 + Q)(z) & S_0(z) \\ S_1(z) & (S_1 + Q)(z) \end{pmatrix} \quad \text{with } z = e^{2\pi i k},$$

which leads to $\chi_{\max}^B = \mathfrak{m}(Q)$.

Example: Newman polynomials and choices

For the class of $\{0, 1\}$ -polynomials, also known as *Newman polynomials*, one has $R = 0$. The associated Fourier matrix is

$$B(k) = \begin{pmatrix} (S_0 + Q)(z) & S_0(z) \\ S_1(z) & (S_1 + Q)(z) \end{pmatrix} \quad \text{with } z = e^{2\pi i k},$$

which leads to $\chi_{\max}^B = \mathfrak{m}(Q)$.

If either S_0 or S_1 is zero, $M_\varrho = B(0)$ is a triangular matrix, and cannot be primitive. This can be avoided if there are at least two columns with a coincidence.

Example: Newman polynomials and choices

For the class of $\{0, 1\}$ -polynomials, also known as *Newman polynomials*, one has $R = 0$. The associated Fourier matrix is

$$B(k) = \begin{pmatrix} (S_0 + Q)(z) & S_0(z) \\ S_1(z) & (S_1 + Q)(z) \end{pmatrix} \quad \text{with } z = e^{2\pi i k},$$

which leads to $\chi_{\max}^B = \mathfrak{m}(Q)$.

If either S_0 or S_1 is zero, $M_\varrho = B(0)$ is a triangular matrix, and cannot be primitive. This can be avoided if there are at least two columns with a coincidence.

If there is only one such column, one can still construct a primitive substitution by recalling that $\mathfrak{m}(-q) = \mathfrak{m}(q)$, so one only needs to exchange the two bijective column types.

As a concrete example, consider $q(z) = 1 + z^2$. The two standard choices

$$\varrho_q : \begin{cases} 0 \mapsto 000 \\ 1 \mapsto 101 \end{cases} \quad \text{and} \quad \varrho_{q'} : \begin{cases} 0 \mapsto 010 \\ 1 \mapsto 111 \end{cases}$$

both give non-primitive substitutions; in fact, their substitution matrices are not even irreducible. However,

$$\varrho_{-q} : \begin{cases} 0 \mapsto 101 \\ 1 \mapsto 000 \end{cases} \quad \text{and} \quad \varrho_{-q'} : \begin{cases} 0 \mapsto 111 \\ 1 \mapsto 010 \end{cases}$$

are both primitive and aperiodic, and have $\chi_{\max}^B = \mathfrak{m}(q)$. One can see in this example that replacing q by $-q$ really means an exchange of w_0 and w_1 in the definition of ϱ .

Lehmer's polynomial, morphism, and dynamical analogue

Lehmer's polynomial, morphism, and dynamical analogue

$$\ell_L(z) = 1 + z - z^3 - z^4 - z^5 - z^6 - z^7 + z^9 + z^{10}$$

Lehmer's polynomial, morphism, and dynamical analogue

$$\ell_{\mathbf{L}}(z) = 1 + z - z^3 - z^4 - z^5 - z^6 - z^7 + z^9 + z^{10}$$

Recall further that $\ell_{\mathbf{L}}$ is the polynomial with the smallest known positive logarithmic Mahler measure, $\mathfrak{m}(\ell_{\mathbf{L}}) \approx \log(1.176281)$. Here,

$$\varrho_{\ell_{\mathbf{L}}} : \begin{cases} 0 \mapsto 00111111000 \\ 1 \mapsto 11100000011 \end{cases}$$

is one of the eight substitutions that correspond to the polynomial $\ell_{\mathbf{L}}$.

Lehmer's polynomial, morphism, and dynamical analogue

$$\ell_L(z) = 1 + z - z^3 - z^4 - z^5 - z^6 - z^7 + z^9 + z^{10}$$

Recall further that ℓ_L is the polynomial with the smallest known positive logarithmic Mahler measure, $\mathfrak{m}(\ell_L) \approx \log(1.176281)$. Here,

$$\varrho_{\ell_L} : \begin{cases} 0 \mapsto 00111111000 \\ 1 \mapsto 11100000011 \end{cases}$$

is one of the eight substitutions that correspond to the polynomial ℓ_L .

Lehmer's problem (dynamical analogue). Does there exist a primitive, 2-automatic sequence ϱ with maximal Lyapunov exponent

$$0 < \chi_{\max}^B < \mathfrak{m}(\ell_L) \approx \log(1.176280818) ?$$

End (fin)